

Do not take these products at the same time.

Alka-Seltzer Plus® Severe Cold + Flu Day

Drug Facts Active ingredients (in each packet) **Purposes** Acetaminophen 500 mg......Pain reliever/fever reducer Dextromethorphan hydrobromide 20 mg......Cough suppressant Guaifenesin 400 mg.....Expectorant

Phenylephrine hydrochloride 10 mg......Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold or flu:
 - minor aches and pains headache
 - sore throat
- cough
- nasal congestion
- · sinus congestion and pressure
- · helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- · with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening blisters
 - rash
- hives

- facial swelling
- asthma (wheezing)shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough with excessive phlegm (mucus)



Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- · redness or swelling is present
- new symptoms occur
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.
- · nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- take every 4 hours; do not exceed 6 packets in 24 hours or as directed by a doctor
- adults and children 12 years and over: dissolve contents of one packet in 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- children under 12 years: do not use

Other information

- each packet contains: potassium 5 mg and sodium 6 mg
- store at room temperature

Inactive ingredients acesulfame potassium, anhydrous citric acid, compressible sugar, D&C yellow #10, dental-type silica, FD&C red #40, flavors, maltodextrin, povidone, pregelatinized starch, silicon dioxide, sodium citrate, stearic acid, sucralose, tartaric acid, tribasic calcium phosphate

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)



Alka-Seltzer Plus® Severe Cold + Flu Night

Drug Facts

Uses

- temporarily relieves these symptoms due to a cold or flu:
 - headache
- minor aches and pains
- cough
- sore throat
- nasal congestion
- sinus congestion and pressure
- runny nose
- sneezing
- · temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- - sters rash
- hives
- facial swelling asthma (wheezing) shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressureglaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema



Ask a doctor or pharmacist before use if you are

- · taking the blood thinning drug warfarin
- · taking sedatives or tranquilizers

When using this product

- · do not exceed recommended dosage
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- · fever gets worse or lasts more than 3 days
- · redness or swelling is present
- · new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- take every 4 hours; do not exceed 5 packets in 24 hours or as directed by a doctor
- adults and children 12 years and over: dissolve contents of one packet in 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- children under 12 years: do not use

Other information

- each packet contains: potassium 5 mg and sodium 5 mg
- store at room temperature

Inactive ingredients acesulfame potassium, anhydrous citric acid, compressible sugar, D&C yellow #10, dental-type silica, FD&C red #40, flavors, pregelatinized starch, sodium citrate, sucralose, tartaric acid, tribasic calcium phosphate

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)