

Drug Facts	
Active ingredient	Purpose
<i>(in each chew</i> ) Magnesium hydroxide USP 1200 i	mgSaline laxative
<b>Use</b> relieves occasional constipation (irregularity). This product usually produces bowel movement in $\frac{1}{2}$ to 6 hours.	
<b>Warnings</b> <b>Do not use</b> if you have ever had a or any of its ingredients	an allergic reaction to this product
<ul> <li>Ask a doctor before use if you have</li> <li>kidney disease</li> <li>a magnesium-restricted diet</li> <li>stomach pain, nausea, or vomiting</li> <li>a sudden change in bowel habits that lasts over 14 days</li> </ul>	
Ask a doctor or pharmacist before use if you are taking a prescription drug. This product may interact with certain prescription drugs.	
<ul> <li>Stop use and ask a doctor if</li> <li>you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.</li> <li>you need to use a laxative for more than 1 week</li> </ul>	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
<ul> <li>Directions</li> <li>drink a full glass (8 oz) of liquid with each dose.</li> <li>dose may be taken as a single daily dose, in divided doses, or as directed by a doctor.</li> <li>do not exceed the maximum recommended daily dose in a 24 hour period</li> </ul>	
adults and children 12 years of age and older	2 to 4 chews
children 6 to under 12 years of age	1 to 2 chews
children 4 to under 6 years of age	1 chew
children under 4 years of age	ask a doctor
<ul> <li>Other information</li> <li>each chew contains: magnesium 500 mg, sodium 1 mg</li> <li>store between 20° to 25° C (68° to 77° F). Avoid excessive heat and humidity.</li> <li>Do not use if pouch is torn or open</li> </ul> Inactive ingredients corn syrup, FD&C red no. 40 aluminum	
lake, flavors, glycerin, hydrogenate sucrose, water.	ed coconut oil, soy lecithin,
Questions or comments? 1-800-647-2529	