



Alka-Seltzer Plus Severe Cold PowerFast Fizz (Orange Zest Flavor, Effervescent Tablets)

Drug Facts

Active ingredients (in each tablet)

Aspirin 325 mg (NSAID)*.....Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg.....Antihistamine
Phenylephrine bitartrate 7.8 mgNasal decongestant
*nonsteroidal anti-inflammatory drug

Purposes

Uses

- temporarily relieves these symptoms due to a cold:
 - minor aches and pains
 - nasal congestion
 - sinus congestion and pressure
 - temporarily reduces fever
 - headache
 - sneezing
 - runny nose
 - sore throat

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children.

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have
 - asthma
 - thyroid disease
 - difficulty in urination due to enlargement of the prostate gland
 - a breathing problem such as emphysema or chronic bronchitis
 - a sodium-restricted diet
 - diabetes
 - glaucoma



Alka-Seltzer Plus Severe Cold PowerFast Fizz (Orange Zest Flavor, Effervescent Tablets)

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for
 - gout • diabetes • arthritis
 - taking sedatives or tranquilizers
-

When using this product

- **do not exceed recommended dosage**
 - excitability may occur, especially in children
 - you may get drowsy
 - avoid alcoholic drinks
 - alcohol, sedatives, and tranquilizers may increase drowsiness
 - be careful when driving a motor vehicle or operating machinery
-

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
 - you experience any of the following signs of stomach bleeding
 - feel faint • vomit blood • have bloody or black stools
 - have stomach pain that does not get better
 - pain or nasal congestion gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - ringing in the ears or a loss of hearing occurs
 - nervousness, dizziness, or sleeplessness occurs
-

If pregnant or breast-feeding, ask a health professional before use.

It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 4 hours. Do not exceed 8 tablets in 24 hours or as directed by a doctor.
 - children under 12 years: do not use
-

Other information

- **each tablet contains:** potassium 80 mg; sodium 356 mg
 - store at room temperature. Avoid excessive heat.
-

Inactive ingredients anhydrous citric acid, calcium silicate, dimethicone, FD&C red #40, FD&C yellow #6, flavor, mannitol, potassium bicarbonate, povidone, sodium bicarbonate, sucralose

Questions or comments? 1-800-986-0369 (Mon – Fri 9AM – 5PM EST)