

Drug Facts Active ingredients (in each tablet) Purpose Aspirin 500 mg (NSAID)*.....Pain reliever Caffeine 65 mg.....Alertness aid/pain reliever aid *nonsteroidal anti-inflammatory drug Uses · for the temporary relief of minor aches and pains associated with a handover helps restore mental alertness or wakefulness when experiencing fatigue or drowsiness associated with a hangover for temporary relief of headaches, body aches, and pain alone Warnings Reve's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction which may include: hives facial swelling asthma (wheezing) shock Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeinecontaining medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat. • For occasional use only. Do not use for more than 2 days for a hangover unless directed by a doctor. Not intended for use as a substitute for sleep. If fatigue or drowsiness persists or continues to recur. consult a doctor. Do not use · if you are allergic to aspirin or any other pain reliever/fever reducer if you have ever had an allergic reaction to this product or any of its ingredients Ask a doctor before use if stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn • • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease •

- you are taking a diuretic
- vou have asthma •
- you have a sodium-restricted diet



 diabetes go 	or out ● arthritis	
5		
top use and ask		
	ion occurs. Seek medical h	
	any of the following signs of /omit blood • have bloody	
	h pain that does not get bet	
	pre than 10 days or gets wor	
redness or swell		
	rs or a loss of hearing occur	S
new symptoms of	occur	
pregnant or bre	ast-feeding, ask a health p	rofessional hefore
	lly important not to use as	
	ncy unless definitely direct	
	may cause problems in the	
complications du		
	of children. In case of ove	erdose, get medic
	oison Control Center right a	
Directions		
fully dissolve 2 t	ablets in 4 ounces of water l	pefore taking
Adults 18 years	2 tablets every 6 hours	do not exceed
and over	as needed, or as	8 tablets in 24
	directed by a doctor	hours
		1
	2 toblete even Charme	do not over a -
	2 tablets every 6 hours	do not exceed
	as needed, or as	5 tablets in 24
nd over	as needed, or as	5 tablets in 24
nd over	as needed, or as directed by a doctor	5 tablets in 24
hildren under 8 years	as needed, or as directed by a doctor consult a doctor	5 tablets in 24
nd over hildren under 8 years D ther informatio	as needed, or as directed by a doctor consult a doctor	5 tablets in 24
Children under 8 years Other information 9 each tablet con	as needed, or as directed by a doctor consult a doctor on ntains: potassium 115 mg	5 tablets in 24
nd over Children under 8 years Other informationer and tablet com each tablet com	as needed, or as directed by a doctor consult a doctor on tains: potassium 115 mg tains: sodium 371 mg	5 tablets in 24 hours
nd over children under 8 years Other informatio each tablet con each tablet con Phenylketonuri	as needed, or as directed by a doctor consult a doctor on ntains: potassium 115 mg ntains: sodium 371 mg ics: Contains Phenylalanine	5 tablets in 24 hours 5.4 mg per tablet
Children under 8 years Children under 8 years Chher information each tablet con each tablet con Phenylketonuri store at room te	as needed, or as directed by a doctor consult a doctor on ntains: potassium 115 mg ntains: sodium 371 mg its: Contains Phenylalanine mperature. Avoid excessive	5 tablets in 24 hours 5.4 mg per tablet heat and humidit
 each tablet con Phenylketonuri store at room ter 	as needed, or as directed by a doctor consult a doctor consult a doctor on ntains: potassium 115 mg itains: sodium 371 mg ics: Contains Phenylalanine mperature. Avoid excessive ents anhydrous citric acid,	5 tablets in 24 hours 5.4 mg per tablet heat and humidit aspartame,
And over Children under 8 years Other information each tablet com each tablet com Phenylketonuri store at room ter mactive ingredio dimethicone, docus	as needed, or as directed by a doctor consult a doctor consult a doctor on ntains: potassium 115 mg itains: sodium 371 mg ics: Contains Phenylalanine mperature. Avoid excessive ents anhydrous citric acid, sate sodium, FD&C red No.	5 tablets in 24 hours 5.4 mg per tablet heat and humidit aspartame, 40, FD&C yellow
Children under 8 years Children under 8 years Chher information each tablet com each tablet com Phenylketonuri store at room tel nactive ingredio limethicone, docus lo. 6, flavor, man	as needed, or as directed by a doctor consult a doctor on ntains: potassium 115 mg itains: sodium 371 mg ics: Contains Phenylalanine mperature. Avoid excessive ents anhydrous citric acid, sate sodium, FD&C red No. nitol, potassium bicarbonate,	5 tablets in 24 hours 5.4 mg per tablet heat and humidit aspartame, 40, FD&C yellow
Children under 8 years Other informatic each tablet com each tablet com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com com	as needed, or as directed by a doctor consult a doctor consult a doctor datains: potassium 115 mg datains: sodium 371 mg ics: Contains Phenylalanine mperature. Avoid excessive ents anhydrous citric acid, sate sodium, FD&C red No. bitol, potassium bicarbonate, bicarbonate, sucralose	5 tablets in 24 hours 5.4 mg per tablet heat and humidit aspartame, 40, FD&C yellow povidone, sodiur
Children under 8 years Other informatic each tablet con each tablet con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con con	as needed, or as directed by a doctor consult a doctor on ntains: potassium 115 mg itains: sodium 371 mg ics: Contains Phenylalanine mperature. Avoid excessive ents anhydrous citric acid, sate sodium, FD&C red No. nitol, potassium bicarbonate,	5 tablets in 24 hours 5.4 mg per table heat and humid aspartame, 40, FD&C yellow povidone, sodiu