

Do not take these products at the same time.

Alka-Seltzer Plus<sup>®</sup> Sinus Congestion & Pain Maximum Strength Day PowerMax<sup>®</sup> Gels

Acetaminophen 325 mgPa Dextromethorphan hydrobromide 10 mg	ain reliever/fever reducer
Dextromethorphan hydrobromide 10 mg	
Phenylephrine hydrochloride 5 mg	
Uses	
<ul> <li>temporarily relieves these symptoms due to</li> </ul>	<ul> <li>cough</li> </ul>
Warnings	
Liver warning: This product contains acetan damage may occur if you take • more than 4,000 mg of acetaminophen in 2 • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while the Allergy alert: Acetaminophen may cause set	24 hours 1 using this product
allergic reactions. Symptoms may include: • skin reddening • blisters • rash • • facial swelling • asthma (wheezing) • If a skin or general allergic reaction occurs, st help right away. <b>Sore throat warning:</b> If sore throat is severe 2 days, is accompanied or followed by fever, or vomiting, consult a doctor promptly.	top use and seek medical , persists for more than
<ul> <li>Do not use</li> <li>with any other drug containing acetaminoph nonprescription). If you are not sure wheth acetaminophen, ask a doctor or pharmacis</li> <li>if you are now taking a prescription monoar (MAOI) (certain drugs for depression, psyci conditions, or Parkinson's disease), or for the MAOI drug. If you do not know if your p an MAOI, ask a doctor or pharmacist befor</li> <li>if you have ever had an allergic reaction to ingredients</li> <li>in children under 12 years of age</li> </ul>	er a drug contains t. mine oxidase inhibitor hiatric, or emotional 2 weeks after stopping prescription drug contains re taking this product.
Ask a doctor before use if you have • liver disease • heart disease • thyroid disease • diabetes • cough that occurs with excessive phlegm (in difficulty in urination due to enlargement of • persistent or chronic cough such as occurs or emphysema	mucus) the prostate gland



## When using this product do not exceed recommended dosage

### Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- · nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not take more than the recommended dose
- do not take the Day and Night products at the same time; wait 4 hours after the last Night dose before starting Day product.
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 6 capsules in 12 hours or as directed by a doctor.
- children under 12 years: do not use

## Other information

 store at room temperature. Avoid excessive heat above 40°C (104°F).

*Inactive ingredients* FD&C yellow #6, ferric oxide, gelatin, glycerin, polyethylene glycol, potassium aluminum silicate, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM -5PM EST)



Alka-Seltzer Plus<sup>®</sup> Sinus Congestion & Pain Maximum Strength Night PowerMax<sup>®</sup> Gels

Drug Facts
Active ingredients (in each capsule) Purposes
Acetaminophen 325 mgPain reliever/fever reducer Dextromethorphan hydrobromide 10 mgPoin reliever/fever reducer Doxylamine succinate 6.25 mgAntihistamine Phenylephrine hydrochloride 5 mgNasal decongestant
Uses <ul> <li>temporarily relieves these symptoms due to a cold or flu:</li> <li>minor aches and pains</li> <li>headache</li> <li>cough</li> <li>sore throat</li> <li>nasal congestion</li> <li>sneezing</li> <li>runny nose</li> <li>sinus congestion and pressure</li> <li>temporarily reduces fever</li> </ul>
<ul> <li>Warnings</li> <li>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</li> <li>more than 4,000 mg of acetaminophen in 24 hours</li> <li>with other drugs containing acetaminophen</li> <li>3 or more alcoholic drinks every day while using this product</li> <li>Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:</li> <li>skin reddening • blisters • rash • hives</li> <li>facial swelling • asthma (wheezing) • shock</li> <li>If a skin or general allergic reaction occurs, stop use and seek medical help right away.</li> <li>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</li> <li>Do not use to sedate children.</li> </ul>
<ul> <li>Do not use</li> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> <li>if you have ever had an allergic reaction to this product or any of its ingredients</li> <li>in children under 12 years of age</li> </ul>
Ask a doctor before use if you have liver disease



### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

#### When using this product

- · do not exceed recommended dosage
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

#### Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- · fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attentior is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not take more than the recommended dose
- do not take the Day and Night products at the same time; wait 4 hours after the last Day dose before starting Night product.
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 4 capsules in 12 hours or as directed by a doctor.
- children under 12 years: do not use

### Other information

 store at room temperature. Avoid excessive heat above 40°C (104°F).

**Inactive ingredients** D&C yellow No. 10, FD&C blue No. 1, gelatin, glycerin, polyethylene glycol, potassium aluminum silicate, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

**Questions or comments? 1-800-986-0369** (Mon-Fri 9AM - 5PM EST)