

Alka-Seltzer Plus Cold & Flu FizzyChews (Chewable Tablets, Orange Flavor)

Drug Facts

Active ingredients (in each chewable tablet) cetaminophen 162.5 mg......Pain reliever/fever reducer

Chlorpheniramine maleate 2 mg......Antihistamine Dextromethorphan hydrobromide 5 mg......Cough suppressant

sneezina

- temporarily relieves these symptoms due to a cold or flu:
 - headache minor aches and pains
 - runny nose

temporarily reduces fever

sore throat

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening blisters rash hives
- facial swelling asthma (wheezing) shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children.

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease glaucoma
- cough that occurs with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- · taking sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase the drowsiness
- be careful when driving a motor vehicle or operating machinery



Alka-Seltzer Plus Cold & Flu FizzyChews (Chewable Tablets, Orange Flavor)

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 tablets every 4 hours.
 Do not exceed 12 tablets in 24 hours or as directed by a doctor.
- do not swallow tablets whole. Chew or crush tablets completely before swallowing.
- children under 12 years: do not use

Other information

- each tablet contains: sodium 17 mg
- store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients anhydrous citric acid, betadex, carboxymethylcellulose, colloidal silicon dioxide, ethylcellulose, flavors, magnesium stearate, mannitol, microcrystalline cellulose, polyethylene, sodium bicarbonate, sodium carbonate, sodium starch glycolate, stearic acid, sucralose, xylitol

Questions or comments? 1-800-986-0369 (Mon - Fri 9AM - 5PM EST)