

Drug Facts

Active ingredient (in each chewable tablet) Purpose

Loratadine 5 mg.....Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

■ chew or crush tablets completely before swallowing.	
adults and children 6 years and over	chew 2 tablets daily; not more than 2 chewable tablets in 24 hours
children 2 to under 6 years of age	chew 1 tablet daily; not more than 1 chewable tablet in 24 hours
children under 2 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- phenylketonurics: contains phenylalanine 1.4 mg per chewable tablet
- safety sealed: do not use if the individual blister unit imprinted with Children's Claritin® is open or torn
- store between 20° to 25°C (68° to 77°F)

Inactive ingredients

aspartame, citric acid anhydrous, colloidal silicon dioxide, flavor, magnesium stearate, mannitol, microcrystalline cellulose, sodium starch glycolate, stearic acid

Questions or comments?

1-800-CLARITIN (1-800-252-7484)