



MiraSOFT Stool Softener Liquid Gels

Drug Facts

Active ingredient (in each softgel)

Docusate sodium 100 mg.....Stool softener laxative

Purpose**Use**

- for relief of occasional constipation (irregularity)
- this product generally produces a bowel movement in 12 to 72 hours

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

Drug Interaction Precaution: Do not take this product if you are presently taking mineral oil, unless directed by a doctor.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take softgels with a full glass (8 oz) of water

adults and children 12 years and older	take 1 to 3 softgels daily or as directed by a doctor. This dose may be taken as a single daily dose or in divided doses.
children 6 to under 12 years	take 1 softgel daily or as directed by a doctor
children under 6 years of age	ask a doctor

Other information

- **each softgel contains:** sodium 5 mg
- store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients gelatin, glycerin, pharmaceutical ink, polyethylene glycol, purified water, sorbitol sorbitan solution. Softgels are imprinted with edible pharmaceutical ink.

Questions or comments? 1-800-647-2529

(Mon – Fri 9AM – 5PM EST)