Bayer HealthCare
Consumer Care

Aspirin Regimen
Bayer® Chewable Low Dose Aspirin 81 mg
Orange Flavored

Drug Facts

Active ingredient (in each chewable tablet)	Purpose
Aspirin 81 mg (NSAID)*….....	Pain reliever

*nonsteroidal anti-inflammatory drug

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor
- ask your doctor about other uses for Bayer 81 mg Orange Chewable Aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
- hives • facial swelling • asthma (wheezing) • shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use
- if you are allergic to aspirin or any other pain reliever/fever reducer
- this product for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for
- gout • diabetes • arthritis

Stop use and ask a doctor if
- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint • vomit blood • have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Bayer HealthCare
Consumer Care

Aspirin Regimen
Bayer® Chewable Low Dose Aspirin 81 mg
Orange Flavored

Directions
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information
- save carton for full directions and warnings
- Store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients colloidal silicon dioxide, corn starch, dextrose, FD&C yellow #6 aluminum lake, flavor, microcrystalline cellulose, saccharin sodium

Questions or comments? **1-800-331-4536** (Mon-Fri 9AM – 5PM EST)