



MiraLAX[®] – Bottles

Drug Facts	
Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g (cap filled to line).....	Purpose Osmotic Laxative
Use <ul style="list-style-type: none">relieves occasional constipation (irregularity)generally produces a bowel movement in 1 to 3 days	
Warnings <p>Allergy alert: Do not use if you are allergic to polyethylene glycol</p> <p>Do not use if you have kidney disease, except under the advice and supervision of a doctor</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none">nausea, vomiting or abdominal paina sudden change in bowel habits that lasts over 2 weeksirritable bowel syndrome <p>Ask a doctor or pharmacist before use if you are taking a prescription drug</p> <p>When using this product you may have loose, watery, more frequent stools</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none">you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.you get diarrheayou need to use a laxative for longer than 1 week <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	
Directions <ul style="list-style-type: none">do not take more than directed unless advised by your doctorthe bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)adults and children 17 years of age and older:<ul style="list-style-type: none">use once a dayfill to top of white section in cap which is marked to indicate the correct dose (17g)stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drinkdo not combine with starch-based thickeners used for difficulty swallowingensure that the powder is fully dissolved before drinkingdo not drink if there are any clumpsdo not use more than 7 dayschildren 16 years of age or under: ask a doctor	
Other information <ul style="list-style-type: none">store at 20° - 25°C (68° - 77°F)tamper-evident: do not use if foil seal under cap, printed with “SEALED for YOUR PROTECTION” is missing, open or broken	
Inactive ingredients none	
Questions or comments? 1-800-MiraLAX (1-800-647-2529)	