



**Alka-Seltzer Plus®
Day & Night Cold & Flu Formula Liquid Gels
Co-Packaged Product**

Do not take these products at the same time.

Alka-Seltzer Plus® Day Non-Drowsy Cold & Flu Formula Liquid Gels

Drug Facts	
Active ingredients (in each capsule)	Purposes
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant
Phenylephrine hydrochloride 5 mg.....	Nasal decongestant
Uses	
<ul style="list-style-type: none"> temporarily relieves these symptoms due to a cold or flu: <ul style="list-style-type: none"> minor aches and pains headache cough sore throat nasal and sinus congestion temporarily reduces fever 	
Warnings	
<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> more than 10 capsules in 24 hours, which is the maximum daily amount for this product with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Allergy Alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:</p> <ul style="list-style-type: none"> skin reddening blisters rash hives facial swelling asthma (wheezing) shock <p>If a skin or general allergic reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>	
Do not use	
<ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. if you have ever had an allergic reaction to this product or any of its ingredients in children under 12 years of age 	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> liver disease heart disease high blood pressure thyroid disease diabetes cough with excessive phlegm (mucus) difficulty in urination due to enlargement of the prostate gland persistent or chronic cough such as occurs with smoking, asthma, or emphysema 	
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin	
When using this product do not exceed recommended dosage	

Alka-Seltzer Plus® Day & Night Cold & Flu Formula Liquid Gels Co-Packaged Product

<p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> • pain, cough, or nasal congestion gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. • nervousness, dizziness, or sleeplessness occurs <hr/> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>
<p>Directions</p> <ul style="list-style-type: none"> • do not take more than the recommended dose • adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor. • children under 12 years: do not use
<p>Other information</p> <ul style="list-style-type: none"> • store at room temperature. Avoid excessive heat above 40°C (104°F).
<p>Inactive ingredients FD&C red #40, FD&C yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide</p>
<p>Questions or comments? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)</p>

Alka-Seltzer Plus® Night Cold & Flu Formula Liquid Gels

Drug Facts	
Active ingredients (in each capsule)	Purposes
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine
Phenylephrine hydrochloride 5 mg.....	Nasal decongestant
Uses	
<ul style="list-style-type: none"> • temporarily relieves these symptoms due to a cold or flu: <ul style="list-style-type: none"> • minor aches and pains • nasal and sinus congestion • runny nose • headache • cough • sneezing • sore throat • temporarily reduces fever 	
Warnings	
<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> • more than 10 capsules in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product 	



**Alka-Seltzer Plus®
Day & Night Cold & Flu Formula Liquid Gels
Co-Packaged Product**

Allergy Alert: acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening • blisters • rash • hives
- facial swelling • asthma (wheezing) • shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease • heart disease • high blood pressure
- thyroid disease • diabetes • glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.



**Alka-Seltzer Plus®
Day & Night Cold & Flu Formula Liquid Gels
Co-Packaged Product**

Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years: do not use

Other information

- store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients FD&C blue #1, D&C yellow #10, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)