

## Drug Facts

### Active ingredient (in each spray)

Azelastine HCl 205.5 mcg (equivalent to 187.6 mcg azelastine)..... Antihistamine

### Purpose

### Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- nasal congestion
- runny nose
- sneezing
- itchy nose

### Warnings

Only for use in the nose. Do not spray in eyes or mouth.

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients

#### Ask a doctor before use if you

- have had recent nose ulcers or nose surgery
- have had a nose injury that has not healed

#### When using this product

- drowsiness may occur
  - avoid alcoholic drinks
  - alcohol, sedatives, and tranquilizers may increase drowsiness
  - be careful when driving a motor vehicle or operating machinery
- you may get a bitter taste in your mouth. To help avoid this, tilt your head downward while spraying.
- nasal discomfort or sneezing may occur right after use
- do not share this bottle with anyone else as this may spread germs

#### Stop use and ask a doctor if

- an allergic reaction, such as a skin rash, to this product occurs
- you have severe or frequent nosebleeds

**if pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- Read the User Guide for how to:
  - prime the bottle before first use
  - prime bottle again if not used for 3 or more days
  - use the spray
  - clean the spray nozzle if it gets clogged

|  |   |
|--|---|
| adults and children 12 years and older | This product may be used either once or twice a day: <ul style="list-style-type: none"><li>■ <b>once daily:</b> use 2 sprays in each nostril; <b>OR</b></li><li>■ <b>twice daily:</b> use 1 or 2 sprays in each nostril every 12 hours</li><li>■ <b>do not use more than 4 sprays</b> in each nostril in a 24 hour period</li></ul> |
| children 6 years to 11 years           | <ul style="list-style-type: none"><li>■ an adult should supervise use</li><li>■ 1 spray in each nostril every 12 hours</li><li>■ <b>do not use more than 2 sprays</b> in each nostril in a 24 hour period</li></ul>   |
| children under 6 years                 | do not use  |

## ***Drug Facts*** (continued)

### ***Other information***

- store between 20° to 25°C (68° to 77°F). Protect from freezing.
- keep this carton and the enclosed User Guide for important information
- do not use if tape printed with “sealed for your protection” on top and bottom carton flaps is not intact

### ***Inactive ingredients***

benzalkonium chloride, edetate disodium, hypromellose, purified water, sodium citrate, sorbitol, sucralose

### ***Questions or comments?***

1-800-317-2165

(Mon–Fri, 9AM–5PM EST)

Bayer and the Bayer Cross are registered trademarks of Bayer.

The Astepro mark is owned by Viartis. Used under license.

© 2021 Bayer.

Dist. by:

Bayer HealthCare LLC

Whippany, NJ 07981

Patents: [patents.livewell.bayer.com](https://patents.livewell.bayer.com)

Product of Germany.

C00011462

**Bayer**

**INDOOR & OUTDOOR**

**Allergy Relief**