



Coricidin® HBP Maximum Strength Cold & Flu Day & Night Liquid Gels

The following statement should appear above the Drug Facts box:
Do not take these products at the same time.

The product name should appear outside the Drug Facts box:
Coricidin HBP® Maximum Strength Cold & Flu Day Liquid Gels

Drug Facts	
Active ingredients (in each capsule)	Purposes
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Guaifenesin 200 mg.....	Expectorant
Uses	
<ul style="list-style-type: none"> • temporarily relieves these symptoms due to a cold or flu: <ul style="list-style-type: none"> • minor aches and pains • headache • sore throat • helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive • temporarily reduces fever 	
Warnings	
<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> • more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product <p>Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:</p> <ul style="list-style-type: none"> • skin reddening • blisters • rash • hives • facial swelling • asthma (wheezing) • shock <p>If a skin or general allergic reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p>	
Do not use	
<ul style="list-style-type: none"> • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you have ever had an allergic reaction to this product or any of its ingredients • in children under 12 years of age 	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> • liver disease • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema • cough that is accompanied by excessive phlegm (mucus) 	
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> • pain or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. 	



Coricidin[®] HBP Maximum Strength Cold & Flu Day & Night Liquid Gels

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- do not take the Day and Night products at the same time; wait 4 hours after taking the last Night dose before taking the Day product.
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 6 capsules in 12 hours or as directed by a doctor.
- children under 12 years of age: do not use

Other information

- store at 15°-25°C (59°-77°F)

Inactive ingredients FD&C yellow # 6, gelatin, glycerin, iron oxide black, lecithin, medium-chain triglycerides, mica, polyethylene glycol, potassium aluminum silicate, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

Questions or comments? 1-800-317-2165 (Mon-Fri 9AM - 5PM EST)



Coricidin® HBP Maximum Strength Cold & Flu Day & Night Liquid Gels

The product name should appear outside the Drug Facts box:
Coricidin® HBP Maximum Strength Cold & Flu Night Liquid Gels

Drug Facts	
Active ingredients (in each capsule)	Purposes
Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine
Uses	
<ul style="list-style-type: none"> • temporarily relieves these symptoms due to a cold or flu: <ul style="list-style-type: none"> • minor aches and pains • headache • cough • sore throat • runny nose • sneezing • temporarily reduces fever 	
Warnings	
<p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> • more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product <p>Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:</p> <ul style="list-style-type: none"> • skin reddening • blisters • rash • hives • facial swelling • asthma (wheezing) • shock <p>If a skin or general allergic reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use to sedate children</p>	
Do not use	
<ul style="list-style-type: none"> • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. • if you have ever had an allergic reaction to this product or any of its ingredients • in children under 12 years of age 	



Coricidin® HBP Maximum Strength Cold & Flu Day & Night Liquid Gels

Ask a doctor before use if you have

- liver disease • glaucoma
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- may cause marked drowsiness
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- do not take the Day and Night products at the same time; wait 4 hours after taking the last Day dose before taking the Night product.
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 4 capsules in 12 hours or as directed by a doctor.
- children under 12 years of age: do not use

Other information

- store at 15°-25°C (59°-77°F)

Inactive ingredients D&C yellow No.10, FD&C blue No.1, gelatin, glycerin, lecithin, medium-chain triglycerides, polyethylene glycol, potassium aluminum silicate, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, soybean oil, titanium dioxide

Questions or comments? 1-800-317-2165 (Mon-Fri 9AM - 5PM EST)



Coricidin® HBP Maximum Strength Cold & Flu Day & Night Liquid Gels

PDP Statement of Identity

Coricidin® HBP Maximum Strength Cold & Flu Day Liquid Gels:

Acetaminophen / Pain reliever-fever reducer
Guaifenesin /Expectorant

Coricidin® HBP Maximum Strength Cold & Flu Night Liquid Gels:

Acetaminophen / Pain Reliever-Fever Reducer
Dextromethorphan Hydrobromide / Cough Suppressant
Doxylamine Succinate / Antihistamine

Abbreviations can be used if space constraints on PDP to read as below:

Acetaminophen / Pain Reliever-Fever Reducer
Dextromethorphan HBr / Cough Suppressant
Doxylamine Succinate / Antihistamine

Contact Information-Preferred format:

Questions or comments? 1-800-317-2165 (Mon-Fri 9AM - 5PM EST)

If there are space constrains, the following abbreviated formats may be used.

Questions or comments? 1-800-317-2165 (Mon-Fri 9 - 5 EST)

Or

Questions? 1-800-317-2165

The following statement must appear prominently on outer carton and in bold type:

Maximum Strength Per 4 Hour Dose