



# Coricidin® HBP

## Maximum Strength Cold, Flu & Chest Congestion Liquid Gels

### Drug Facts

| Active ingredients (in each capsule) | Purposes                    |
|--------------------------------------|-----------------------------|
| Acetaminophen 325 mg.....            | Pain reliever/fever reducer |
| Guaifenesin 200 mg.....              | Expectorant                 |

### Uses

- temporarily relieves these symptoms due to a cold or flu:
  - minor aches and pains
  - headache
  - sore throat
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

### Ask a doctor before use if you have

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that is accompanied by excessive phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

### Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.



**Coricidin® HBP**  
**Maximum Strength Cold, Flu & Chest Congestion Liquid Gels**

|  |
|--|
| <b>Directions</b> <ul style="list-style-type: none"><li>• do not take more than the recommended dose</li><li>• adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.</li><li>• children under 12 years of age: do not use</li></ul> |
| <b>Other information</b> <ul style="list-style-type: none"><li>• store at 15°-25°C (59°-77°F)</li></ul>  |
| <b>Inactive ingredients</b> FD&C yellow # 6, gelatin, glycerin, iron oxide black, lecithin, medium-chain triglycerides, mica, polyethylene glycol, potassium aluminum silicate, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide                              |
| <b>Questions or comments? 1-800-317-2165</b> (Mon-Fri 9AM - 5PM EST)   |

**Contact Information - Preferred format:**

**Questions or comments? 1-800-317-2165** (Mon-Fri 9AM - 5PM EST)

If there are space constraints, the following abbreviated formats may be used.

**Questions or comments? 1-800-317-2165** (Mon-Fri 9 - 5 EST)

Or **Questions? 1-800-317-2165**

**PDP Statement of Identity:**

**Acetaminophen** / Pain Reliever-Fever Reducer  
Guaifenesin /Expectorant

The following statement must appear prominently on outer carton and in bold type:

**Maximum Strength Per 4 Hour Dose**